


Specification

Guanidine Thiocyanate for molecular biology

A1107

Solubility:	1420 g/L (H ₂ O)
Physical Description:	Solid
Product Code:	A1107
Product Name:	Guanidine Thiocyanate for molecular biology
Specifications:	DNases/RNases/Proteases: not detectable Assay (titr.): min. 99 % pH (1 M): 4.7 - 7.0 Water (K.F.): max. 1.0 % Ammonium: max. 0.1 % Fe: max. 0.0005 % A (1 cm/6 M in water HPLC grade) 280 nm: max. 0.6 300 nm: max. 0.1
Hazard pictograms	
WGK:	2
Storage:	RT
Signal Word:	Attention
GHS Symbols:	GHS07
H Phrases:	EUH032 H302+H312+H332 H412
P Phrases:	P273

AppliChem GmbH

Ottoweg 4 • D-64291 Darmstadt • Phone +49 6151 9357 0 • Fax +49 6151 9357 11 • info.de@itwreagents.com • www.itwreagents.com
 CEO Joan Roget • Commerzbank Darmstadt • Bank 508 800 50 • Account 0186989900 IBAN DE24 5088 0050 0186 9899 00 • Swiftcode
 DRESDEFF508 • Finanzamt Darmstadt 07 228 16476 • Register court Darmstadt HRB Nr. 7340

Specification

Guanidine Thiocyanate for molecular biology

A1107

	P302+P352
Molecular Formula:	CH ₆ N ₃ SCN
M:	118.16 g/mol
CAS:	593-84-0
EINECS:	209-812-1
CS:	29252900
Index Nr.:	615-004-00-3
Comment Reference (1) describes the original protocol for the guanidine thiocyanate (GuaSCN) / β-mercaptoethanol methode for the isolation of non-degraded RNA from RNase-rich tissues (e. g. pancreas). There are many variations from this method published (e. g. ref. 3, 4).As an alternative to the toxic formaldehyde, guanidine thiocyanate may be applied for Northern blotting (5). A final concentration of 20 mM is sufficient to denature RNA in the agarose gel (5).	
Bibliography (1)Chirgwin, J.M. <i>et al.</i> (1979) <i>Biochemistry</i> 18 , 5294-5299227Isolation of biologically active ribonucleic acid from sources enriched in ribonuclease. (2)MacDonald, R.J. <i>et al.</i> (1987) <i>Methods Enzymol.</i> 152 , 219-227Isolation of RNA using guanidinium salts. (3)Lizardi, P.M. (1983) <i>Methods Enzymol.</i> 96 , 24-38Methods of the preparation of messenger RNA. (4)Chomczynski, P. & Sacchi, N. (1987) <i>Anal. Biochem.</i> 162 , 156-159Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. (5)Goda, S.K. & Minton, N.P. (1995) <i>Nucleic Acids. Res.</i> 23 , 3357-3358A simple procedure for gel electrophoresis and Northern blotting of RNA.	

AppliChem GmbH

Ottoweg 4 • D-64291 Darmstadt • Phone +49 6151 9357 0 • Fax +49 6151 9357 11 • info.de@itwreagents.com • www.itwreagents.com
 CEO Joan Roget • Commerzbank Darmstadt • Bank 508 800 50 • Account 0186989900 IBAN DE24 5088 0050 0186 9899 00 • Swiftcode DRESDEFF508 • Finanzamt Darmstadt 07 228 16476 • Register court Darmstadt HRB Nr. 7340