



**PRODUCT CODE: 251550**

**Benedict's Reagent qualitative for clinical diagnosis**  
for determination of glucose in urine

---

TARIC 3822 00 00 00

**PHYSICAL DATA:** liquid, Clear, Dark blue, Miscible with water • D 20/4 1,153 •

**INSTRUCTIONS FOR USE:** 5 ml of Benedict's reagent + 10 drops of urine. Boil two minutes and allow to cool. If the solution becomes cloudy, it indicates the presence of glucose.

**BIBLIOGRAPHY:** Manual Técnico de Análisis Clínicos, Eduardo Suarez, 9 edic., 861-862 •

**HAZARDOUS:**

H: H412 •

P: P273 • P501 •

**WEIGHT/VOLUME INFORMATION:** 1l~1,153 kg      1kg~0,867 l

**COMPOSITION:**

Sodium Carbonate anhydrous	153,6 g
Citric Acid 1-hydrate	70,9 g
Copper(II) Sulphate 5-hydrate	15,6 g
Water (s.q.m)	1 l

**SPECIFICATIONS:**

Identity :  
Identity passes test

**Maximum limit of impurities**

Suitability for determination of glucose in urine passes test

Ed.: 3 . Vig.: 30.08.2017 .

Prod.: 251550