



**PRODUCT CODE: 161429****Butanone, 99.5% (Methylethylketone) for synthesis**

M.= 72,11

CAS [78-93-3]

EINECS 201-159-0

TARIC 2914 12 00 00

FEMA 2170

SYNONYMS: Ethyl Methyl Ketone, MEK, Methyl Ethyl Ketone•

PHYSICAL DATA: liquid, Clear, Colourless, Soluble in water 290 g/l at 20 °C D 20/4 0,805 • M.P.: -86 °C • B.P.: 79,6 °C • pH(300g/l)5,5 • n₂₀/D : 1,379 • Flash P.: -6 °C • Ign. T.:505 °C • Vap. press. (20 °C) 105 hPa • Viscosity 15 °C 0,38 mPa.s • D. M. 20 °C 2,7 Debye • Dielec. constant20 °C 18,5 • Evap. number (DIN 53170) 6,3 • Heat evap. 80 °C 444 KJ/Kg • Satur. conc. 20 °C 310 g/m³ • Expl. limit1,8 %(V)11,5 %(V) •

BIBLIOGRAPHY: Merck Index **12**, 6.149 13, 6.097 Sax **BOV250** • Safety **2** , **606 C** • Kühn-Birett **M 14** • Ullmann **(4.)14** , 193 • Beilstein **1** , **666 I** , **347 II** , **726 III** ,**2270 IV** , **4243** • BRN 741880 • ACS **X216IX200** • F.C.C **8 9** • Directive 88/344/E.C.E.92/115/E.C.E.94/52/EC97/60/EC (27/10/1997) 2009/32/CE • Royal Decree 472/1990 (6/4/1990) , 2667/1998 (11 /12/1998), 1101/2011 (22/7/2011) •

HAZARDOUS: C.E: 606-002-00-3 • RTECS: EL 6475000 • LD50 oral rat 2.737 mg/kg • LC50 rat 23500mg/m³ / 8h • LD50 skn rbt 6.480 mg/kg • VLA-EC 300 ppm900 mg/m³ VLA-ED 200 ppm600 mg/m³



H: H225 • H319 • EUH066 • H336 •

P: P210 • P233 • P240 • P241 • P242 • P501 • P243 • P261 • P264 • P271 • P280 • P303+P361+P353 • P304+P340 • P305+P351+P338 • P312 • P337+P313 • P370+P378 • P403+P233 • P403+P235 • P405 •

TRANSPORT REGULATIONS: UN: 1193 • ADR: 3/II • IMDG: 3/II • IATA: 3/II • PAX: 353 • CAO: 364 • (D/E) •**WEIGHT/VOLUME INFORMATION:** 1l~0,806 kg 1kg~1,241 l

OBSERVATIONS: Product controlled as a drug precursor.

SPECIFICATIONS:

Minimum assay (G.C.)	99,5%
Identity :	
Identity	IR passes test
Density at 20/4	0,804-0,808
Non-volatile matter	0,002 %
Water (H ₂ O)	0,2 %

Ed.: 4 . Vig.: 15.01.2011 .

Prod.: 161429